Printed from NOAH Compendium (https://hub21.community.librios.com). (c) Copyright NOAH Compendium 2026. All Rights Reserved.
Date: Thursday, February 19, 2026 15:24

Virbac Limited

Telephone:01359 243243

Website:www.virbac.com

Email:enquiries@virbac.co.uk

Cortavance0.584mg/ml cutaneous spray solution for dogs (GB only)

Species: Dogs

Therapeutic indication:Pharmaceuticals: Anti-inflammatory preparations: Topical

Active ingredient: Hydrocortisone Aceponate

Product:CORTAVANCE

Product index:CORTAVANCE

Incorporating:

Qualitative and quantitative composition

Hydrocortisone aceponate 0.584 mg/ml

Pharmaceutical form

Cutaneous spray, solution.

Clinical particulars

Target species

Dogs

Indications for use

For symptomatic treatment of inflammatory and pruritic dermatoses in dogs.

For alleviation of clinical signs associated with atopic dermatitis in dogs.

Contra-indications

Do not use on cutaneous ulcers.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients

Special precautions for use

Special precautions for use in animals

Clinical signs of atopic dermatitis such as pruritus and skin inflammation are not specific for this disease and therefore other causes of dermatitis such as ectoparasitic infestations and infections which cause dermatological signs should be ruled out before treatment is started, and underlying causes should be investigated.

In the case of concurrent microbial disease or parasitic infestation, the dog should receive appropriate treatment for such condition.

In the absence of specific information, the use in animal suffering from Cushing's syndrome shall be based on the risk-benefit assessment.

Since glucocorticosteroids are known to slow growth, use in young animals (under 7 months of age) shall be based on the risk-benefit assessment and subject to regular clinical evaluations.

Total body surface treated should not exceed approximately 1/3 of the dog's surface corresponding for example to a treatment of two flanks from the spine to the mammary chains including the shoulders and the thighs. Otherwise, use only according to the risk-benefit assessment of the responsible veterinary surgeon and subject the dog to regular clinical evaluations as further described below.

Care should be taken to avoid spraying into the eyes of the animal.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

The active substance is potentially pharmacologically active at high doses of exposure.

The formulation may cause eye irritation following accidental ocular contact.

The formulation is flammable.

Wash hands after use. Avoid contact with eyes.

To avoid skin contact, recently treated animals should not be handled until the application site is dry.

To avoid inhalation of the product, apply the spray in a well-ventilated area.

Do not spray on naked flame or any incandescent material.

Do not smoke while handling the veterinary medicinal product.

Replace the bottle in the outer carton and in a safe place out of the sight and the reach of children immediately after use.

In case of accidental skin contact, avoid hand-to-mouth contact and wash the exposed area immediately with water.

In case of accidental eye contact, rinse with abundant quantities of water.

If eye irritation persists, seek medical advice.

In case of accidental ingestion, especially by children, seek medical advice immediately and show the leaflet or the label to the physician.

Other precautions

The solvent in this product may stain certain materials including painted, varnished or other household surfaces or furnishings. Allow the application site to dry before permitting contact with such materials.

Adverse Reactions

Transient local reactions at the application site (erythema and/or pruritus) can occur in very rare cases.

Use during pregnancy or lactation

The safety of the veterinary medicinal product has not been established during pregnancy and lactation. Systemic absorption of hydrocortisone aceponate being negligible, it is unlikely for teratogenic, foetotoxic, maternotoxic effects to happen at the recommended dosage in dogs.

Use only according to the risk-benefit assessment by the responsible veterinarian.

Interactions

In the absence of information, it is recommended not to apply other topical preparations simultaneously on the same lesions.

Amounts to be administered and administration route

Cutaneous use.

Before administration, screw the pump spray on the bottle.

The veterinary medicinal product is then applied by activating the pump spray, from a distance of about 10 cm of the area to be treated.

The recommended dosage is 1.52 µg of hydrocortisone aceponate/cm2 of affected skin per day. This dosage can be achieved with two pump spray activations over a surface to be treated equivalent to a square of 10 cm x 10 cm.

- For treatment of inflammatory and pruritic dermatoses, repeat the treatment daily for 7 consecutive days. In case of conditions requiring an extended treatment, the responsible veterinarian should subject the use of the veterinary medicinal product to the risk-benefit assessment. If signs fail to improve within 7 days, treatment should be re-evaluated by the veterinarian.

- For alleviation of clinical signs associated with atopic dermatitis, repeat the treatment daily for at least 14 and up to 28 consecutive days. An intermediary control by the veterinarian at day 14 should be made to decide if further treatment is needed. The dog should be re-evaluated regularly with regard to HPA suppression or skin atrophy, both being possibly asymptomatic. Any prolonged use of

this product, to control atopy, should be at the benefit risk assessment of the responsible veterinary surgeon. It should take place after a re-evaluation of the diagnosis and also a consideration of the multi-modal treatment plan in the individual animal.

Presented as a volatile spray, this veterinary medicinal product does not require any massage.

Overdose

Tolerance studies of multiple doses were assessed over a period of 14 days in healthy dogs using 3 and 5 times the recommended dosage corresponding to the two flanks, from the spine to the mammary chains including the shoulder and the thighs (1/3 of the dog's body surface area). These resulted in a reduced capacity for production of cortisol that is fully reversible within 7 to 9 weeks after the end of treatment.

In 12 dogs suffering from atopic dermatitis, after topical application once a day at the recommended therapeutic dosage for 28 to 70 (n=2) consecutive days, no noticeable effect on the systemic cortisol level was observed.

Withdrawal periods

Not applicable.

Pharmacological particulars

Pharmacotherapeutic group: Corticosteroids, dermatological preparations.

ATCvet code: QD07AC16.

Pharmacodynamic properties

The veterinary medicinal product contains the active substance hydrocortisone aceponate.

Hydrocortisone aceponate is a dermocorticoid with a potent intrinsic glucocorticoid activity which means a relief of both inflammation and pruritus leading to a quick improvement of skin lesions observed in case of inflammatory and pruritic dermatosis. In case of atopic dermatitis, improvement will be slower.

Pharmacokinetic properties

HCA belongs to the diesters class of the glucocorticosteroids.

The diesters are lipophilic components ensuring an enhanced penetration into the skin associated to a low plasma availability. HCA thus accumulates in the dog's skin allowing local efficacy at low dosage. The diesters are transformed inside the skin structures. This transformation is responsible for the potency of the therapeutic class. In laboratory animals, hydrocortisone aceponate is eliminated the same way as hydrocortisone (other name for endogenous cortisol) through urine and faeces.

Topical application of diesters results in high therapeutic index: high local activity with reduced systemic secondary effects.

Pharmaceutical particulars

Excipients

Propylene glycol methyl ether

Major incompatibilities

None known.

Shelf life

Shelf-life of the veterinary medicinal product as packaged for sale: 2 years.

Shelf-life after first opening the immediate packaging: 6 months.

Special precautions for storage

None

Nature and composition of immediate packaging

Box containing a polyethylene terephtalate (PET) or high density polyethylene (HDPE) bottle filled with 31 ml or 76 ml of solution, closed with an aluminium screw cap or a white plastic screw cap and a pump spray.

Carton box with a PET bottle of 31ml

Carton box with a PET bottle of 76ml

Carton box with a HDPE bottle of 31ml

Carton box with a HDPE bottle of 76ml

Not all pack sizes may be marketed.

Disposal

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Marketing Authorisation Number

MARKETING AUTHORISATION HOLDER

Virbac

Premiere Avenue

2065M – L I D

06516 Carros Cedex

France

GB: Vm 05653/5001

Significant changes

Date of the first authorisation or date of renewal

09.01.2007

Date of revision of the text

July 2021

Any other information

Keep treated animals away from fires, other sources of heat and surfaces likely to be affected by the excipient (propylene glycol methyl ether) for at least 30 minutes following spraying or until the fur is totally dry.

Legal category

Legal category:POM-V

GTIN

GTIN description: Cortavance 76ml

GTIN:3597133039498