

NOAH Compendium

Printed from NOAH Compendium (<https://hub21.community.librios.com>). (c) Copyright NOAH Compendium 2026. All Rights Reserved.

Date: Tuesday, February 17, 2026 15:20

[Norbrook Laboratories Ltd](#)

Telephone: 01536 741147

Website: www.norbrook.com

Email: gbenquiries@norbrook.co.uk

Lincoject 10 % w/v Solution for Injection

Species: Cats, Dogs, Pigs

Therapeutic indication: Pharmaceuticals: Antimicrobials: Injections

Active ingredient: Lincomycin

Product: Lincoject 10% w/v Solution for Injection

Product index: Lincoject Injection

Pig - meat: 3 days

Incorporating:

Qualitative and quantitative composition

Active substance:

Lincomycin	10.0% w/v [100 mg/ml]
------------	-----------------------

(as Lincomycin hydrochloride)	11.34% w/v [113.4 mg/ml]
-------------------------------	--------------------------

Excipients:

Benzyl Alcohol	0.9% w/v [9mg/ml]
----------------	-------------------

For a full list of excipients, see section Pharmaceutical Particulars

Pharmaceutical form

Solution for Injection.

A clear, colourless solution for injection.

Clinical particulars

Target species

Pigs, Dogs, Cats

Indications for use, specifying the target species

In dogs and cats, Lincoject is recommended for the treatment of infections caused by Lincomycin susceptible Gram-positive organisms, particularly streptococci and staphylococci and certain Gram-negative anaerobic bacteria (e.g. *Bacteroides*, *Fusobacterium* species).

Lincoject is recommended for treatment of the following conditions in dogs and cats: Tonsillitis, laryngitis and other respiratory tract infections; abscesses, infected wounds and purulent dermatitis; septicaemia.

In pigs, Lincoject is indicated for the treatment of infections caused by Lincomycin susceptible Gram-positive organisms (e.g. staphylococci, streptococci), certain Gram-negative anaerobic organisms (e.g. *Serpulina* (*Treponema*) *hyodysenteriae*, *Bacteroids*, *Fusobacterium* and *Mycoplasma* species).

Lincoject is recommended for treatment of the following conditions in pigs: Swine dysentery, enzootic or mycoplasmal pneumonia, septic arthritis, foot abscesses.

Contraindications

The use of Lincoject is not recommended in species other than the cat, dog and pig. Lincosamides may cause fatal enterocolitis in horses, rabbits and rodents and diarrhoea and reduced milk production in cattle.

Lincoject should not be given to animals with a known pre-existing monilial infection.

Not to be used in animals hypersensitive to Lincomycin

Special Warnings for each target species

When treating groups of pigs, the use of a multiple dose syringe is recommended. To refill the syringe, the use of a draw-off needle is recommended to avoid excessive broaching of the stopper.

Special precautions for use

Special precautions for use in animals

Do not exceed the recommended dosage.

Practice aseptic injection techniques.

Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local epidemiological information.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Avoid contact with the product. In case of accidental eye or skin contact, wash off the affected area thoroughly with water.

Adverse reactions (frequency and seriousness)

The intramuscular administration of Lincoject to pigs at higher levels than recommended may result in diarrhoea and loose stools.

Use during pregnancy, lactation or lay

Not applicable.

Interaction with other medicinal products and other forms of interaction

Not compatible with penicillin and erythromycin if used concurrently.

Amounts to be administered and administration route

Lincoject is indicated for intravenous and intramuscular administration to dogs and cats and intramuscular administration to pigs.

CATS AND DOGS:

Intramuscularly: 22 mg/kg of bodyweight once daily or 11 mg/kg every

12 hours.

Intravenously: 11 to 22 mg/kg one or two times per day by slow intravenous injection.

PIGS:

Intramuscularly: 4.5 to 11 mg/kg of bodyweight once daily.

To ensure the correct dosage, bodyweight should be determined as accurately as possible to avoid underdosing.

Overdose (symptoms, emergency procedures, antidotes), if necessary

The intramuscular administration of Lincoject to pigs at higher levels than recommended may result in diarrhoea and loose stools.

Withdrawal period

Animals must not be slaughtered for human consumption during treatment.

Pigs (Meat): 3 days

Pharmacological particulars

Pharmacotherapeutic group: Antibiotic

ATC Vet Code: QJ01FF02

Pharmacodynamic properties

The mode of action is inhibition of protein synthesis at the ribosomal 50S subunit level. Lincomycin is primarily excreted in the faeces as both parent compound and metabolites with large biliary contribution. After administration of the recommended single dosage, faecal excretion accounts for 38% and urinary excretion for 49% of the total dose. The antibiotic is transported in polymorphonuclear leucocytes to the site of infection, an event that may explain the efficient penetration and targeting of organisms in tissues difficult to reach by diffusion in the presence of purulent material.

Pharmaceutical particulars

List of excipients

- Benzyl Alcohol
- Water for Injections

Incompatibilities

None known

Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 2 Years.

Shelf life after first opening the immediate packaging: 28 Days.

Special precautions for storage

Do not store above 25°C. Protect from light.

Following withdrawal of the first dose use the product within 28 days. Discard unused product.

Nature and composition of immediate packaging

50 ml and 100 ml Type I clear colourless glass vials, complete with bromobutyl bungs and aluminium caps.

Not all pack sizes may be marketed.

Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

NOAH Compendium

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Marketing Authorisation Holder (if different from distributor)

Norbrook Laboratories Limited, Station Works, Camlough Road, Newry, Co. Down, BT35 6JP Northern Ireland

Marketing Authorisation Number

Vm 02000/4183

Significant changes

Date of the first authorisation or date of renewal

20th November 2001

Date of revision of the text

April 2022

Any other information

Nil.

Legal category

Legal category:POM-V

GTIN

GTIN description:Lincoject 50ml

GTIN:5023534003021

GTIN description:Lincoject 100ml

GTIN:5023534003038