

Dosage and administration

For external use only (spot-on use)


The recommended minimum doses are 10 mg/kg bodyweight imidacloprid and 1.0 mg/kg bodyweight moxidectin, equivalent to 0.1 ml/kg bodyweight of the veterinary medicinal product. The treatment schedule should be based on individual veterinary diagnosis and on the local epidemiological situation.

Administer in accordance with the following table:

Dosage	Weight of animal (kg)	Volume (ml)
Prinocate 40mg/4mg spot-on solution for small cats and ferrets	<4	0.4
Prinocate 80mg/8mg spot-on solution for large cats	4-8	0.8
Prinocate 40mg/10mg spot-on solution for small dogs	<4	0.4
Prinocate 100mg/25mg spot-on solution for medium dogs	>4-10	1
Prinocate 250mg/62.5mg spot-on solution for large dogs	>10-25	2.5
Prinocate 400mg/100mg spot-on solution for extra-large dogs	>25-40	4

Method of administration:

1. Remove one pipette from the package. Hold applicator pipette in an upright position, twist and pull cap off.
2. Turn the cap around and place the other end of cap back on pipette. Push and twist the cap to break seal, and then remove the cap from the pipette.
3. With the dog/cat standing still, part the coat between the shoulder blades until the skin is visible. The product should only be applied to undamaged skin. Place the tip of the pipette on the skin and squeeze firmly several times to empty its contents completely and directly onto the skin in one spot. Avoid contact between the veterinary medicinal product and your fingers.



Use during pregnancy, lactation: The safety of the veterinary medicinal product has not been established during pregnancy and lactation. Laboratory studies with either imidacloprid or moxidectin in rats and rabbits have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects. Use only according to the benefit-risk assessment by the responsible veterinarian.