

NOAH Compendium

Printed from NOAH Compendium (<https://hub21.community.librios.com>). (c) Copyright NOAH Compendium 2026. All Rights Reserved.

Date: Monday, February 23, 2026 16:00

[Elanco UK AH Limited](#)

Telephone:01256 353131 (24 hr response) Veterinary Technical Services

Website:www.my.elanco.com/en_gb

Email:elancouk@elanco.com

Veraflox 25 mg/ml oral suspension for cats

Species:Cats

Therapeutic indication:Pharmaceuticals: Antimicrobials: Oral preparations: Others

Active ingredient:Pradofloxacin

Product:Veraflox 25 mg/ml oral suspension for cats

Product index:Veraflox 25 mg/ml oral suspension for cats

Incorporating:

Qualitative and quantitative composition

Each ml contains:

Active substance:

Pradofloxacin 25 mg

Excipients:

Qualitative composition of excipients and other constituents	Quantitative composition if that information is es the veterinary medicinal product
Sorbic acid (E200)	2mg
Amberlite IRP 64	
Ascorbic acid	
Xanthan gum	
Propylene glycol	
Vanilla flavour	
Purified water	

NOAH Compendium

Yellowish to beige suspension.

Pharmaceutical form

Oral suspension

Yellowish to beige suspension

Clinical particulars

Target species

Cats

Indications for use, specifying the target species

Treatment of:

- acute infections of the upper respiratory tract caused by strains of *Pasteurella multocida*, *Escherichia coli* and the *Staphylococcus intermedius* group (including *S. pseudintermedius*).
- wound infections and abscesses caused by strains of *Staphylococcus intermedius* group (including *S. pseudintermedius*) and *Pasteurella multocida*.

Contraindications

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

Do not use in kittens less than 6 weeks.

Pradofloxacin has no effects on the developing cartilage of kittens 6 weeks of age and older. Do not use in cats with persisting articular cartilage lesions, since lesions may worsen during treatment with fluoroquinolones.

Do not use in cats with central nervous system (CNS) disorders, such as epilepsy, as fluoroquinolones could potentially cause seizures in predisposed animals.

Do not use in cats during pregnancy and lactation (see section below: Use during pregnancy, lactation or lay)

Special warnings for each target species

Cross-resistance has been shown between pradofloxacin and other fluoroquinolones. Use of pradofloxacin should be carefully considered when susceptibility testing has shown resistance to fluoroquinolones because its effectiveness may be reduced.

Special precautions for use

Special precautions for safe use in the target species

NOAH Compendium

Use of the product should be based on identification and susceptibility testing of the target pathogen(s). If this is not possible, therapy should be based on epidemiological information and knowledge of susceptibility of the target pathogens at local/regional level.

Use of the product should be in accordance with official, national and regional antimicrobial policies.

An antibiotic with a lower risk of antimicrobial resistance selection (lower AMEG category) should be used for first line treatment where susceptibility testing suggests the likely efficacy of this approach. Narrow spectrum antibiotic therapy with a lower risk of antimicrobial resistance selection should be used for first line treatment where susceptibility testing suggests the likely efficacy of this approach.

Pradofloxacin may increase sensitivity of the skin to sunlight. During treatment, animals should therefore not be exposed to excessive sunlight.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

People with known hypersensitivity to quinolones should avoid any contact with the veterinary medicinal product.

Avoid skin and eye contact with the veterinary medicinal product. Wash hands after use.

In case of accidental contact with the eyes, wash immediately with water.

In case of contact with the skin, rinse off with water.

Do not eat, drink or smoke while handling the veterinary medicinal product.

In case of accidental ingestion, seek medical advice and show the package leaflet or the label to the physician.

Adverse reactions (frequency and seriousness)

Cats.

Rare (1 to 10 animals / 10,000 animals treated):	Digestive tract disorder (e.g.
---	--------------------------------

Use during pregnancy, lactation or lay

The safety of this veterinary medicinal product has not been established during pregnancy and lactation.

Pregnancy:

Do not use during the whole or part of pregnancy. Laboratory studies in rats have shown evidence of pradofloxacin induced eye malformations at foetotoxic and maternotoxic dosages.

NOAH Compendium

Lactation:

Do not use during lactation since there are no data on pradofloxacin in kittens aged less than 6 weeks. Fluoroquinolones are known to cross the placenta and to be distributed into milk.

Fertility:

Pradofloxacin has been shown to have no effects on fertility in breeding animals.

Interaction with other medicinal products and other forms of interaction

Concurrent administration with metal cations such as those contained in antacids or sucralfate made with magnesium hydroxide or aluminium hydroxide, or multivitamins containing iron or zinc, and dairy products containing calcium, has been reported to decrease the bioavailability of fluoroquinolones. Therefore, the veterinary medicinal product should not be administered concurrently with antacids, sucralfate, multivitamins or dairy products, as absorption of the veterinary medicinal product may be decreased.

Further, fluoroquinolones should not be used in combination with non-steroidal anti-inflammatory drugs (NSAIDs) in animals with a history of seizures because of potential pharmacodynamic interactions in the CNS. The combination of fluoroquinolones with theophylline could increase the plasma levels of theophylline by altering its metabolism and thus should be avoided. The combined use of fluoroquinolones with digoxin should also be avoided because of potentially increased oral bioavailability of digoxin.

Amounts to be administered and administration route

Oral use.

The recommended dose is 5 mg/kg bodyweight of pradofloxacin once daily. To ensure a correct dosage, body weight should be determined as accurately as possible. Due to the graduation of the syringe the resulting dose range is 5 to 7.5 mg/kg bodyweight according to the following table:

Dosing Table

Bodyweight of Cat (kg)	Dose of Oral suspension to be given (ml)
> 0.67 - 1	0.2
>1 - 1.5	0.3
>1.5 - 2	0.4
>2 - 2.5	0.5
>2.5 - 3	0.6
>3 - 3.5	0.7
>3.5 - 4	0.8

NOAH Compendium

>4 - 5	1
>5 - 6	1.2
>6 - 7	1.4
>7 - 8	1.6
>8 - 9	1.8
>9 - 10	2

Duration of treatment.


The duration of the treatment depends on the nature and severity of the infection and on the response to treatment. For most infections the following treatment courses will be sufficient:

Indication	Duration of treatment
Wound infections and abscesses	7
Acute infections of the upper respiratory tract	5


The treatment should be reconsidered if no improvement of the clinical condition is observed within 3 days after starting the treatment.

Method of administration


To facilitate accurate dosing, the 15 ml bottle of Veraflox oral suspension is supplied with a 3 ml oral dosing syringe (graduation: 0.1 to 2 ml).



Shake well before use.



Draw out the equivalent dosage into the syringe.



Administer directly into the mouth.

In order to avoid cross contamination, the same syringe should not be used for different animals. Thus, one syringe should only be used for one animal. After administration, the syringe should be cleaned with tap water and stored in the carton box together with the veterinary medicinal product.

Overdose (symptoms, emergency procedures, antidotes), if necessary

NOAH Compendium

No specific antidotes for pradofloxacin (or other fluoroquinolones) are known, therefore, in case of overdose, symptomatic treatment should be given.

Intermittent vomiting was observed after repeated oral administration of 1.6 times the maximum recommended dose.

Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance Not applicable.

Withdrawal period(s)

Not applicable

Pharmacological particulars

ATCvet code: QJ01MA97

Pharmacodynamic properties

Mode of Action

The primary mode of action of the fluoroquinolones involves interaction with enzymes essential for major DNA functions such as replication, transcription and recombination. The primary targets for pradofloxacin are the bacterial DNA gyrase and topoisomerase IV enzymes. Reversible association between pradofloxacin and DNA gyrase or DNA topoisomerase IV in the target bacteria results in inhibition of these enzymes and rapid death of the bacterial cell. The rapidity and extent of bacterial killing are directly proportional to the drug concentration.

Antibacterial Spectrum

Although pradofloxacin has *in-vitro* activity against a wide range of Gram-positive and Gram-negative organisms, including anaerobic bacteria, this veterinary medicinal product should only be used for the approved indications (see section: Indications for use, specifying the target species) and in accordance with the prudent use recommendations in section: special precautions for use of this SPC

MIC-Data

Bacterial Species	Number of strains	MIC50
<i>Pasteurella multocida</i> - respiratory tract infections (RTI)1	64	0.008
<i>Pasteurella multocida</i> – wound infections2	42	0.008
<i>Escherichia coli</i> - respiratory tract infections (RTI)1	22	0.015
<i>Staphylococcus intermedius</i> group (including <i>S. pseudintermedius</i>) - respiratory tract infections (RTI)1	25	0.12

NOAH Compendium

Staphylococcus intermedius group (including *S. pseudintermedius*) – wound infections² 20 0.03

1 Data collected between 2017–2018 2 Data collected between 2021–2022 The bacteria were isolated from clinical cases in Belgium, Czech Republic, France, Germany, Hungary, Italy, the Netherlands, Poland, Spain Sweden, Switzerland and UK.

Clinical breakpoints established by CLSI in 2024 (7th edition) for pradofloxacin in cats for respiratory tract infections are:

Organism	Minimum Inhibitory Concentration breakpoints of pradofloxacin (mcg/	
	Susceptible	intermediate
<i>E.coli</i>	≤0.25	0.5-1
<i>S.pseudintermedius</i>	≤0.25	0.5-1

Types and Mechanisms of Resistance

Resistance to fluoroquinolones has been reported to arise from five sources, (i) point mutations in the genes encoding for DNA gyrase and/or topoisomerase IV leading to alterations of the respective enzyme, (ii) alterations of drug permeability in Gram-negative bacteria, (iii) efflux mechanisms, (iv) plasmid mediated resistance and (v) gyrase protecting proteins. All mechanisms lead to a reduced susceptibility of the bacteria to fluoroquinolones. Cross-resistance within the fluoroquinolones class of antimicrobials is common.

Pharmacokinetic particulars

In laboratory studies the bioavailability of pradofloxacin was reduced in fed cats compared to fasted animals. However in clinical studies feeding did not reveal any impact on the treatment effect.

After oral administration of the veterinary medicinal product to cats at the recommended therapeutic dose, absorption of pradofloxacin is rapid, reaching peak concentrations of 2.1 mg/l within 1 hour. The bioavailability of the veterinary medicinal product is at least 60%. Repeated dosing shows no impact on the pharmacokinetic profile, (accumulation index = 1.2). *In vitro* plasma protein binding is low (30%). The high volume of distribution (Vd) >4 l/kg body weight indicates good tissue penetration. Pradofloxacin is eliminated from serum with a terminal half-life of 7 hours. The major elimination pathway in cats is glucuronidation. Pradofloxacin is cleared from the body at 0.28 l/h/kg.

Pharmaceutical particulars

Major incompatibilities

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 3 years

NOAH Compendium

Shelf life after first opening the immediate packaging: 3 months

Special precautions for storage

Store in original container

Keep the bottle tightly closed

Nature and composition of immediate packaging

Folding carton containing a white high density polyethylene (HDPE) bottle with a polyethylene adapter and a child-resistant closure.

Pack sizes: 15 ml bottle with a 3 ml polypropylene oral dosing syringe (graduation: 0.1 to 2 ml) and 30 ml bottle.

Not all pack sizes may be marketed.

Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater. Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

Marketing Authorisation Holder (if different from distributor)

Bayer Animal Health GmbH

D-51368 Leverkusen

Germany

Marketing Authorisation Number

UK (Great Britain): Vm 04895/5010

UK (Northern Ireland): EU/2/10/107/013-014

Significant changes

Date of the first authorisation or date of renewal

Date of first authorisation: 12/04/2011

Date of last renewal: 07/01/2016

Date of revision of the text

December 2025

Any other information

NOAH Compendium

Legal category

Legal category:POM-V

GTIN

GTIN description:Veraflox 25 mg/ml oral suspension for cats

GTIN:04007221032281